

## Guidance on the use of Pentacel and Pediarix August 2008

Pentacel is a combination vaccine that contains DTaP, IPV and Hib vaccines. Pentacel is supplied as single-dose vials, 5 doses to a package. A single-dose vial of liquid DTaP-IPV vaccine is used to reconstitute a single-dose vial of lyophilized ActHIB vaccine. The vaccine must be kept at refrigerator temperature (35°-46° F) at all times. Pentacel must never be frozen. Vaccine exposed to freezing temperature must not be used.



### IMPORTANT NOTE:

The availability of Pentacel will improve the Hib vaccine supply situation in the United States. However, the availability of Pentacel is not sufficient to reinstate the last (booster) dose of the Hib vaccine series (i.e. the dose administered after the first birthday). Although Pentacel is licensed by FDA for the fourth dose in the DTaP, IPV and Hib series, providers should NOT use it for the fourth dose until there is further improvement in the Hib vaccine supply (anticipated for the last quarter of 2008). **Until the Hib supply improves Pentacel should be used ONLY for the first three doses of the DTaP, IPV, and Hib vaccination series, except as noted below.**

As with all combination vaccines, there are no special rules for Pentacel, except as determined by FDA licensure of the product (i.e., the maximum age for any dose- see below). The schedule, minimum intervals, and minimum ages are determined by the individual components. The recommended schedule for Pentacel is functionally the same as for DTaP and ActHib with doses at 2, 4, 6, and 15 through 18 months of age.

Pentacel can be administered to any child 6 weeks through 4 years of age, without a contraindication to any component, for whom DTaP, IPV, and Hib vaccines are indicated. As stated on the childhood immunization schedule, **a combination vaccine, including Pentacel, may be used whenever any component(s) of the combination is indicated and no other component of the vaccine is contraindicated.** This means that Pentacel can be used when a child needs one or two components, but does not need the others.

Contraindications and precautions for Pentacel are the same as those for DTaP, IPV, and Hib vaccines.

The following minimum ages and intervals are defined for the component vaccines in various ACIP statements, and in particular in Table 1 of the 2006 version of the *General Recommendations on Immunization* (<http://www.cdc.gov/mmwr/PDF/rr/rr5515.pdf>, page 3) and on page 31-32 of the 2006 AAP *Red Book*.

Parameter	Age/interval
Minimum age for any dose	6 weeks
Minimum interval for doses 1 and 2	4 weeks
Minimum age for dose 2	10 weeks
Minimum interval for doses 2 and 3	4 weeks
Minimum age for dose 3	14 weeks
Minimum interval for dose 3 and 4	<b>6 months</b> (determined by DTaP component; minimum interval for dose 3-4 is two months for Hib and four weeks for IPV)
Minimum age for dose 4	<b>12 months</b> (determined by DTaP and Hib components). Note that both the minimum interval AND age must be met for the fourth dose of DTaP or Hib (as Pentacel or any other formulation) to be counted as valid
Maximum age for <b>any</b> dose	4 years, 364 days (i.e., do not administer at age 5 years or older)

Please refer to the tables below for guidance on schedules for Pentacel, Pediarix and the single antigen series for Hep B, Hib, IPV DTaP for healthy children\* during the Hib vaccine shortage.

## Examples of Schedules Using Pentacel and/or Pediarix for Healthy Children\* During the Hib Shortage

The first two tables below provide examples of how to introduce Pentacel in your practice using two different schedules. The second two tables review the schedules for the single antigen and Pediarix series for Hep B, IPV, Hib and DTaP.

### Schedule for Hep B, Hib\*, IPV, and DTaP Using Pentacel for All Doses

Birth	2 mos.	4 mos.	6 mos.	15-18 mos.	4-6 years
Hep B	Hep B		Hep B		
				DTaP	DTaP
					IPV
	Pentacel	Pentacel	Pentacel		

### Schedule for Hep B, Hib\*, IPV, and DTaP Using Pentacel For First Dose Only and Pediarix for Remainder of Doses

Birth	2 mos.	4 mos.	6 mos.	15-18 mos.	4-6 years
Hep B	Hep B				
		Hib	Hib		
				DTaP	DTaP
					IPV
	Pentacel				
		Pediarix	Pediarix		

### Schedule for Hep B, Hib\*, DTaP and IPV Without Pentacel or Pediarix

Birth	2 mos.	4 mos.	6 mos.	15-18 mos.	4-6 years
Hep B	Hep B		Hep B		
	Hib	Hib	Hib		
	DTaP	DTaP	DTaP	DTaP	DTaP
	IPV	IPV	IPV		IPV

### Schedule for Hep B, Hib\*, IPV, and DTaP Using Pediarix Only (No Pentacel)

Birth	2 mos.	4 mos.	6 mos.	15-18 mos.	4-6 years
Hep B					
	Hib	Hib	Hib		
				DTaP	DTaP
					IPV
	Pediarix	Pediarix	Pediarix		

Pentacel contains DTaP, IPV and Hib. Pediarix contains DTaP, IPV, and Hep B

Neither Pentacel nor Pediarix should be used prior to 6 weeks of age. In general ACIP recommends the same brand of DTaP be used for all doses of the series. However, different brands can be used if the providers does not know or have available the brand of DTaP used for prior doses.

\*When supplies are sufficient an additional dose of Hib vaccine (single antigen or as part of a combination vaccine) is recommended for healthy children at 12 through 15 months of age (at least 2 months after the prior dose). Either Pentacel or single antigen Hib vaccine may be used at 12 through 15 months of age for children who are at increased risk of Hib disease or who have not completed a complete primary Hib schedule. If Pentacel is administered at 12 through 15 months of age a dose of DTaP at 15 through 18 months of age is not needed. See *MMWR* 2007;56(No.50):1318-1320 for additional details.

Questions or comments on this document should be directed to the Immunization Services Division, NCIRD, by email at [nipinfo@cdc.gov](mailto:nipinfo@cdc.gov).